

# Continuous Renal Replacement Therapy for Acute Kidney Injuries after Cardiac Surgery - Insights from a Single Cardiovascular Institute

Genta Chikazawa<sup>1,2\*</sup>, Arudo Hiraoka<sup>1</sup>, Tomoya Ohshita<sup>3</sup>, Motoko Kimura<sup>4</sup>, Satoko Ishii<sup>4</sup>, Toshinori Totsugawa<sup>1</sup> and Hidenori Yoshitaka<sup>1</sup>

<sup>1</sup>Department of Cardiovascular Surgery, the Sakakibara Heart Institute of Okayama, Japan

<sup>2</sup>Department of Intensive Care Unit, the Sakakibara Heart Institute of Okayama, Japan

<sup>3</sup>Department of Clinical Engineering, the Sakakibara Heart Institute of Okayama, Japan

<sup>4</sup>Department of Anesthesiology, the Sakakibara Heart Institute of Okayama, Japan

## Abstract

**Background:** Acute Kidney Injury (AKI) is not an unusual clinical complication after cardiovascular surgery. The aim of this study was to investigate the clinical significance of Continuous Renal Replacement Therapy (CRRT) in managing postoperative AKI.

**Material and Methods:** 243 patients who received cardiovascular surgery complicated with postoperative AKI in Cardiovascular Intensive Care (CVICU) were enrolled. Patients were divided into two groups: those requiring CRRT (Group A, n=64) and those not administered CRRT (Group B, n=179).

**Results:** Preoperative backgrounds showed there were more surgically at-risk patients complicated with a poorer renal profile in Group A. Procedure-related data identified Cardio Pulmonary Bypass (CPB) and Aortic Cross Clamp (ACC) times were significantly longer in Group A. Also, in-hospital mortality and hemodialysis transition rates were significantly higher in Group A. Preoperative renal profile and male gender were identified as risk factors for hemodialysis transition post cardiovascular surgery.

**Conclusions:** CRRT is expected to play a vital role in managing AKI post cardiovascular surgery. Further investigation is warranted to clarify the efficacy of CRRT for the improvement of long-term outcomes of patients complicated with postoperative AKI.

**Keywords:** Acute kidney injury; Cardiopulmonary bypass; Continuous renal replacement therapy

**\*Corresponding author:** Genta Chikazawa, Department of Cardiovascular Surgery, the Sakakibara Heart Institute of Okayama, 2-5-1, Nakai-cho, Kita-ku, Okayama, 700-0804, Japan, Tel: +81-86-225-7111, Fax: +81-86-223-5265; E-mail: endochikazawa@yahoo.co.jp

**Received Date:** February 09, 2022

**Accepted Date:** February 15, 2022

**Published Date:** February 22, 2022

**Citation:** Chikazawa G, Hiraoka A, Ohshita T, Kimura M, Ishii S, et al. (2022) Continuous Renal Replacement Therapy for Acute Kidney Injuries after Cardiac Surgery - Insights from a Single Cardiovascular Institute. J Emerg Med Trauma Surg Care 4: 014.

**Copyright:** © 2022 Chikazawa G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**HD:** HemoDialysis

**LOS:** Length of Stay

**BSA:** Body Surface Area

**DM:** Diabetes Mellitus

**COPD:** Chronic Obstructive Pulmonary Disease

**HT:** Hypertension

**Min BT:** Minimum Bladder Temperature

## Abbreviations

**AKI:** Acute Kidney Injury

**CRRT:** Continuous Renal Replacement Therapy

**CPB:** Cardio Pulmonary Bypass

**ACC:** Aortic Cross Clamp

**CVICU:** Cardio Vascular Intensive Care Unit

**CABG:** Coronary Artery Bypass Grafting

**KDIGO:** Kidney Disease Improving Global Outcomes

**ANP:** Atrial Natriuretic Petide

**LVEF:** Left Ventricular Ejection Fraction

**sCre:** serum Creatinine

**eGFR:** estimated Glomerular Filtration Rate

## Introduction

Acute Kidney Injury (AKI) is not an unusual postoperative complication after cardiac surgery and is independently associated with increased mortality and morbidity [1,2]. The incidence of AKI following cardiac surgery varies between 20% and 50%, depending on the diagnostic criteria used to define renal injury [3]. Therapeutic strategies to prevent AKI are imperative in improving clinical outcomes of cardiac surgery as AKI can eventually lead to multi-organ failure if not appropriately treated once AKI is diagnosed [2,4-7]. Continuous Renal Replacement Therapy (CRRT) has been reported to be an effective treatment for AKI following cardiac surgery [8,9]. The primary purpose of this study was to examine the clinical significance of prompt initiation of Continuous Renal Replacement Therapy (CRRT) [8,9] under the diagnosis of AKI following cardiovascular surgery, based on the experience of a single cardiovascular institute.

## Methods

A total of 243 patients who had undergone elective or emergency cardiovascular surgery with Cardiopulmonary Bypass (CPB) complicated with postoperative AKI at our institute's Cardiovascular Intensive Care Unit (CVICU) between 2012 and 2020 were enrolled. Those on dialysis preoperatively or with a functioning renal transplant were excluded. Surgeries included: isolated ischemic heart procedure, such as Coronary Artery Bypass Grafting (CABG); isolated heart valve operation, repair or replacement; thoracic aortic fixation for aneurismal or dissecting disorders; and, combinations of those isolated procedures stated above. AKI was confirmed by the Kidney Disease: Improving Global Outcomes (KDIGO) guidelines which define AKI as an increase in sCr concentration  $\geq 0.3$  mg/dL within 48 hours or an increase in sCr concentration  $\geq 1.5$ X baseline that is known or presumed to have occurred within the prior 7 days, or urine output  $<0.5$  ml/kg/h for 6 hours [10,11]. The initiation of CRRT was routinely introduced as promptly as possible after surgery to prevent and treat volume overload unresponsive to diuretics or systemic administration of Atrial Natriuretic Peptide (ANP), uremic symptoms and signs, exacerbated metabolic acidosis, and hyperkalemia in establishing AKI [12,13]. All the enrolled cases were divided into two groups: those requiring CRRT (Group A, n=64) and those without CRRT (Group B, n=179).

## Statistical Analysis

Continuous data are presented as a median and interquartile range. Normally distributed data were analyzed using 2-tailed *t*-tests, and non-normally distributed data were compared with the Mann-Whitney test, as appropriate. Categorical variables are given as a count and percentage of patients and compared using the  $\chi^2$  test. When any expected frequency was less than 1, or 20% of expected frequencies were less than or equal to 5, Fisher's exact test was used. Clinical outcomes between both groups and the risk factors of chronic hemodialysis transition among the surviving cases were analyzed by multivariate analysis of variance (MANOVA). A *p*-value of  $<0.05$  was considered significant. All data were analyzed using the Statistical Analysis Systems software JMP 12.0 (SAS Institute Inc., Cary, NC, USA).

All of the enrolled patients underwent elective or emergency surgery after written informed consent was obtained from them or their families. In proceeding with this study, approval from our Institutional Review Board was granted.

## Results

Preoperative backgrounds are listed in Table 1. There were significant differences between the groups using the Logistic EuroSCORE for 30-day mortality ( $6.5 \pm 3.6\%$  vs.  $4.3 \pm 3.1\%$ ,  $p<0.001$ ), Left Ventricular Ejection Fraction (LVEF) on preoperative echocardiography ( $50.8 \pm 17.5\%$  vs.  $57.6 \pm 14.0\%$ ,  $p=0.002$ ), Serum Creatinine (sCr) ( $1.73 \pm 0.91$  mg/dl vs.  $1.12 \pm 0.37$  mg/dl,  $p<0.001$ ), or estimated Glomerular Filtration Rate (eGFR) ( $36.3 \pm 18.2$  ml/min vs.  $50.0 \pm 15.6$  ml/min,  $p<0.001$ ). Those in Group A were revealed to be at a surgically higher risk, complicated with reduced Left Ventricular Function (LVEF) and poorer renal profiles. Procedure-related data is shown in Table 2. Although there were no significant differences between the groups with regard to systemic perfusion pressure, minimum bladder temperature during CPB, and intraoperative total amount of urination, the Aortic Cross Clamp (ACC) time ( $143.9 \pm 50.0$  min vs.  $126.6 \pm$

$42.6$  min,  $p=0.009$ ) and CPB time ( $222.6 \pm 86.7$  min vs.  $184.1 \pm 59.0$  min,  $p<0.001$ ) were significantly longer in Group A than in Group B. In addition, the total amount of Blood Transfusion (BTF) ( $1785 \pm 1308$  ml vs.  $1032 \pm 1026$  ml,  $p<0.001$ ) during the entire procedure was much higher in Group A than in Group B. In the present study, both redo and specific type of surgery, or the complexities of more than two isolated procedures, were not significantly associated with the requirement of CRRT for postoperative AKI. Postoperative data is shown in Table 3. Postoperative hospital stay (POH) ( $52.3 \pm 41.4$  days vs.  $30.2 \pm 22.9$  days,  $p<0.001$ ), CVICU stay ( $52.3 \pm 41.4$  days vs.  $30.2 \pm 22.9$  days,  $p<0.001$ ), and intubation time following operation ( $148.0 \pm 253$  hours vs.  $26.7 \pm 37.6$  hours,  $p<0.001$ ) were significantly longer in Group A than in Group B.

Variables	Group A (n = 64)	Group B (n = 179)	P value
Age (year)	$72.5 \pm 10.6$	$73.0 \pm 10.7$	.67
Female sex (n)	21 (33%)	63 (35%)	.76
Logistic Euro-score (%) *mortality	$6.5 \pm 3.6$	$4.3 \pm 3.1$	<001
BSA (m <sup>2</sup> )	$1.58 \pm 0.20$	$1.62 \pm 0.19$	.09
LVEF (%)	$50.8 \pm 17.5$	$57.6 \pm 14.0$	.002
Redo Surgery(%)	13 (20%)	21 (12%)	.22
sCr (mg/dl)	$1.73 \pm 0.91$	$1.12 \pm 0.37$	<001
Hgb (g/dl)	$12.0 \pm 2.9$	$12.7 \pm 1.8$	.009
eGFR (ml/min/1.73m <sup>2</sup> )	$36.3 \pm 18.2$	$50.0 \pm 15.6$	<001
DM(n)	38 (59%)	85 (47%)	.11
COPD(n)	9 (14%)	39 (22%)	.20
HT(n)	31 (48%)	90 (50%)	.88

Table 1: Preoperative Backgrounds.

Variables	Group A (n = 64)	Group B (n = 179)	P value
CPB time (min)	$222.6 \pm 86.7$	$184.1 \pm 59.0$	$P<001$
ACC time (min)	$143.9 \pm 50.0$	$126.6 \pm 42.6$	.009
Perfusion Pressure (mmHg)	$57.7 \pm 11.2$	$55.3 \pm 9.5$	<001
Min BT (degree)	$33.5 \pm 1.1$	$33.6 \pm 1.0$	.48
Urine Output (ml)	$703.6 \pm 1095.9$	$510.6 \pm 517.2$	.07

Table 2: Procedure Related Data.

Variables	Group A (n = 64)	Group B (n = 179)	P value
Hospital LOS (day)	$52.3 \pm 41.4$	$30.2 \pm 22.9$	$P<001$
ICU LOS (day)	$14.2 \pm 14.4$	$4.4 \pm 3.6$	$P<001$
Intubation Time (hour)	$148.0 \pm 253.0$	$26.7 \pm 37.6$	<001
In-hospital Death (%)	12.5% *8 cases	1.7% *3 cases	.001
HD transition Rate (%) *survival case	32% *18 cases	0%	$P<001$

Table 3: Postoperative Data.

It was not unusual that hospital death was markedly higher in Group A ( $12.5\%$  vs.  $26.7 \pm 1.7\%$ ,  $p=0.001$ ).

Regarding the surviving cases at discharge in both groups, the Hemo Dialysis (HD) transition rate was 32% in Group A and 0% in

Group B, which was statistically significant ( $p < 0.001$ ). Risk factors of chronic HD transition are shown in Table 4. While male gender and preoperative sCr or eGFR were revealed significant on univariate analysis, only preoperative sCr was identified on multivariate analysis.

	Univariate Analysis		Multivariate Analysis	
	OR(CI)	P value	Adjusted OR(CI)	P value
Male gender	4.05(1.1-9.6)	0.034		
sCre value	6.91(2.6-26.3)	<001	6.87(1.01-66.1)	049
eGFR value	0.92(0.86-0.97)	<001		

**Table 4:** Risk factors for HD transition.

## Discussion

Postoperative administration of prompt CRRT for AKI following cardiac surgery was reported to be effective in decreasing operative mortality, reducing the CVICU Length Of Stay (LOS) as well as hospital LOS, and reducing the duration of CRRT. The benefits for early introduction of CRRT are considered to be effective in decreasing the chance of life-threatening complications including uremia, acidemia, volume overload, and hyperkalemia. Also, initiation of CRRT as promptly as possible, which balances electrolyte, solute, and fluid levels before exacerbation to a more serious stage can effectively reduce the risk of kidney-specific and non-kidney vital organ injury, compared to a late introduction of CRRT.

Once AKI is confirmed, we routinely initiate CRRT for postoperative AKI which is unresponsive to diuretics and systemic administration of renal-protective agents, such as ANP.

However, there are some negative or controversial studies which are opposed to early initiation of CRRT because it could expose patients to potential harms, such as hemorrhage, bacteremia derived from the placement of the dialysis catheter, and thrombosis, intradialytic hypotension, clearance of trace elements, and hypersensitivity to the extracorporeal circuit or antibiotics, which could add to resource utilization [14]. Thus, the optimal treatment strategy for AKI is not only the prompt initiation of CRRT after the operation but also the careful intraoperative management which can help avoid the need for CRRT introduction.

Maintaining renal perfusion during CPB is key to the prevention of AKI [2]. Typical hemodynamic goals during CPB are to support normal organ function and maintenance of a mean perfusion pressure of 50 to 70 mmHg, and these values are based on experimental studies of minimum blood flow required to deliver adequate oxygenation [15,16]. In the present study, mean perfusion pressure during CPB between the groups was not significantly different, and perfusion pressure settings and determination by our institute perfusionists was considered to be appropriate so as to preserve renal perfusion. Karkouti K. demonstrated BTF in excess of 2 units consistently increased the risk of patients developing AKI [17]. Considering the amount of BTF was significantly higher in Group A than in Group B in our study, secure and complete hemostasis during surgery should be attained in order to prevent the initiation of CRRT for exacerbated AKI. Also, needless to say CBP time under cardiac arrest should be shortened to as little as possible to avoid the use of CRRT because ACC time as well as CPB time were significantly longer in Group A than in Group B. In short, a sophisticated surgical technique is an important factor for the prevention of AKI requiring CRRT following cardiac surgery.

Regarding postoperative management, hemodynamic instability such as low cardiac output is a primary cause of AKI after cardiac surgery [3]. This can occur from multiple factors, including hypovolemia, acute ventricular dysfunction such as perioperative myocardial infarction, prolonged ACC time, persistent hemorrhage, or cardiac tamponade. Prompt diagnosis is crucial, and the main aspects of treatment strategies, including goal-directed fluid management and administration of renal protective agents, such as ANP and dexamethasone [3,13,18,19], can be implemented prior to the initiation of CRRT. In the present study, those in Group A were identified as more surgically high-risk, with worse renal profile, and poorer clinical outcomes, including significantly higher rates of hemodialysis transition and in-hospital mortality. Accordingly, in the real setting of postoperative management, prompt determination to initiate CRRT for AKI unresponsive to those conservative treatments stated above should be performed, especially for more severe and complicated cases.

Our study has several limitations. Data are derived from a small sample size at a single cardiovascular institute. Therefore, our models need to be evaluated at multi-centers to be validated. Further investigation is needed to clarify the long-term outcomes of all the surviving cases at discharge for conclusive clinical significance of CRRT introduction after cardiovascular surgery using CPB [20-22].

## Conclusion

AKI following cardiovascular surgery is not an unusual complication, and a comprehensive approach, including goal-directed fluid management, administration of renal protective agents, and CRRT initiation should be implemented. Of these treatments of choice, CRRT has the potential to improve early clinical outcomes of those cases complicated with AKI. Thus, prompt initiation of CRRT should be considered, especially for more surgically severe and complicated cases.

## References

1. O'Neal JB, Shaw AD, Billings IV (2016) Acute kidney injury following cardiac surgery: Current understanding and future directions. *Critical Care* 20: 187.
2. Chertow GM, Levy EM, Hammermeister KE, Grover F, Daley J (1998) Independent association between acute renal failure and Mortality following cardiac surgery. *Am J Med* 104: 343-348.
3. Chew STH, Hwang NC (2020) Acute kidney Injuries after cardiac surgery: A narrative review of the literature. *Journal of Cardiothoracic and Vascular Anesthesia* 33: 1122-1138.
4. Kumar AB, Suneja M (2011) Cardiopulmonary bypass- associated acute kidney injury. *Anesthesiology* 114: 964-970.
5. Brown JR, Kramer RS, Coca SG, Parikh CR (2010) Duration of acute kidney injury impacts long-term survival after cardiac surgery. *Ann Thorac Surg* 90: 1142-1149.
6. Srivastava V, D'Silva C, Tang A, Sogliani F, Ngaage DL (2012) The impact of major perioperative renal insult on long-term renal function and survival after cardiac surgery. *Interactive Cardiovascular and Thoracic Surgery* 15: 14-17.
7. Coleman MD, Shaefi S, Sladen RN (2011) Preventing acute kidney injury after cardiac surgery. *Current Opinion in Anesthesiology* 24: 70-76.
8. Zou H, Hong Q, XU G (2017) Early versus late initiation of renal replacement therapy impacts mortality in patients with acute kidney injury post cardiac surgery. *Critical Care* 150.
9. Vidal S, Richebe P, Barandon L, Calderon J, Tafer N (2009) Evaluation of continuous veno-venous hemofiltration for the treatment of cardiogenic

- shock in conjunction with acute renal failure after cardiac surgery. *European Journal of Cardio-thoracic Surgery* 36: 572-579.
10. Khwaja A (2012) KDIGO Clinical Practice Guidelines for Acute Kidney Injury. *Nephron Clin Pract* 120: c179-c184.
  11. Machado MN, Nakazone MA, Maia LN (2014) Acute kidney injury based on KDIGO criteria in patients with elevated baseline serum creatinine undergoing cardiac surgery. *Rev Bras Cir Cardiovasc* 29: 299-307.
  12. Parolari A, Pesce LL, Pacini D, Mazzanti V, Salis S, et al. (2012) Risk factors for perioperative acute kidney injury after cardiac surgery: Role of perioperative management. *Ann Thorac Surg* 93: 584-591.
  13. Nigwekar SU, Navaneethan SD, Parikh CR, Hix JK (2009) Atrial natriuretic peptide for management of acute kidney injury: A systematic Review and Meta-analysis. *Clin J Am Soc Nephrol* 4: 261-272.
  14. Gibney RT, Bagshaw SM, Kutsogiannis DJ, Johnston C (2008) When should renal replacement therapy for acute kidney injury be initiated and discontinued? *Blood Purif* 26: 473-484.
  15. Ranucci M, Romitti F, Isgro G, Cotza M, Brozzi S, et al. (2005) Oxygen delivery during cardiopulmonary bypass and acute renal failure after coronary operations. *Ann Thorac Surg* 80: 2213-2220.
  16. Somer FD, Mulholland JW, Bryan MR, Alosio T, Nooten GJV, et al. (2011) O<sub>2</sub> delivery and CO<sub>2</sub> production during cardiopulmonary bypass as determinants of acute kidney injury: Time for a goal-directed perfusion management? *Critical Care* 15: R192.
  17. Karkouti K (2012) Transfusion and risk of acute kidney injury in cardiac surgery. *Br J Anaesth* 109: i29-i38.
  18. Cho JS, Shim JK, Soh S, Kim MK, Kwak YL (2016) Perioperative dexmedetomidine reduces the incidence and severity of acute kidney injury following valvular heart surgery. *Kidney International* 89: 693-700.
  19. Jacob KA, Leaf DE, Dieleman JM, Dijk D, Nierich AP (2015) Intraoperative High-Dose Dexamethasone and Severe AKI after Cardiac Surgery. *J Am Nephrol* 26: 2947-2951.
  20. Fielitz AH, Hasse M, Bellomo R, Calzavacca P, Spura A, et al. (2017) Perioperative hemodynamic instability and fluid overload are associated with increasing acute kidney injury severity and worse outcome after cardiac surgery. *Blood Purif* 43:298-308.
  21. McGuinness SP, Parke RL, Bellomo R, Van Haren FM, Bailey M (2013) Sodium Bicarbonate Infusion to reduce Cardiac Surgery-Associated Acute Kidney Injury: A Phase II Multicenter Double-Blind Randomized Controlled Trial. *Crit Care Med* 41: 1599-1607.
  22. Newland RF, Tully PJ, Baker RA (2013) Hyperthermic perfusion during cardiopulmonary bypass and postoperative temperature are independent predictors of acute kidney injury following cardiac surgery. *Perfusion* 28: 223-231.



Henry Journal of Acupuncture & Traditional Medicine

Henry Journal of Anesthesia & Perioperative Management

Henry Journal of Aquaculture and Technical Development

Henry Journal of Cardiology & Cardiovascular Medicine

Henry Journal of Case Reports & Imaging

Henry Journal of Cell & Molecular Biology

Henry Journal of Tissue Biology & Cytology

Henry Journal of Clinical, Experimental and Cosmetic Dermatology

Henry Journal of Diabetes & Metabolic Syndrome

Henry Journal of Emergency Medicine, Trauma & Surgical Care

Henry Journal of Haematology & Hemotherapy

Henry Journal of Immunology & Immunotherapy

Henry Journal of Nanoscience, Nanomedicine & Nanobiology

Henry Journal of Nutrition & Food Science

Henry Journal of Obesity & Body Weight

Henry Journal of Cellular & Molecular Oncology

Henry Journal of Ophthalmology & Optometry

Henry Journal of Perinatology & Pediatrics

Submit Your Manuscript: <https://www.henrypublishinggroups.com/submit-manuscript/>

Henry Publishing Groups, 41341 Red Birch Dr, Aldie, VA, 20105, USA

Tel: +1 571-275-4480; E-mail: [contact@henrypublishinggroups.org](mailto:contact@henrypublishinggroups.org)

<https://www.henrypublishinggroups.com/>

---